

# Necessary and Sufficient Conditions for Optimal Control of Semilinear Stochastic PDEs

Lukas Wessels

joint work with Wilhelm Stannat

Technische Universität Berlin

*[www.lukaswessels.org](http://www.lukaswessels.org)*

PDE Seminar, Georgia Tech, September 20, 2022



Berlin Mathematical School

**BMS**



# Example

Consider Nagumo equation


$$\partial_t x_t^u = \Delta x_t^u + x_t^u(x_t^u - 1)(a - x_t^u) + u_t, \quad x_0 = x \in L^2(\Lambda),$$

with Neumann boundary conditions, where  $\Lambda \subset \mathbb{R}$  bounded and  $a \in (0, 1)$ .


Introduce control  $u : [0, T] \times \Lambda \rightarrow \mathbb{R}$ . Minimize

$$J(u) = \int_0^T \int_{\Lambda} (x_t^u(\lambda) - x_{\bar{\Lambda}}(t, \lambda))^2 d\lambda dt + \int_{\Lambda} (x_T^u(\lambda) - x^T(\lambda))^2 d\lambda$$


for given reference profiles  $x_{\bar{\Lambda}}, x^T$ .



no control



controlled



applied control

# Motivation

Consider deterministic control problem: Minimize

$$J(u) := \int_0^T \int_{\Lambda} l(x_t^u(\lambda), u_t) d\lambda dt + \int_{\Lambda} h(x_T^u(\lambda)) d\lambda$$

subject to

$$\partial_t x_t^u = \Delta x_t^u + b(x_t^u, u_t).$$

Introduce random input disturbance  $\xi_t(\lambda)$ , i.e.

$$\partial_t x_t^u = \Delta x_t^u + b(x_t^u, u_t + \xi_t).$$

Taylor expansion yields

$$b(x_t^u, u_t + \xi_t) \approx b(x_t^u, u_t) + b_u(x_t^u, u_t)\xi_t.$$

Leads to controlled stochastic PDE

$$dx_t^u = [\Delta x_t^u + b(x_t^u, u_t)]dt + b_u(x_t^u, u_t)dW_t,$$

where  $(W_t)_{t \geq 0}$  is an (infinite-dimensional) Brownian motion.

Minimize

$$J(u) := \mathbb{E} \left[ \int_0^T \int_{\Lambda} l(x_t^u(\lambda), u_t) d\lambda dt + \int_{\Lambda} h(x_T^u(\lambda)) d\lambda \right]$$

subject to

$$\begin{cases} dx_t^u = [\Delta x_t^u + b(x_t^u, u_t)] dt + \sigma(x_t^u, u_t) dW_t, & t \in [0, T] \\ x_0^u = x_0 \in L^2(\Lambda), \end{cases}$$

where

- $l, h$  Nemytskii operators of (at most) quadratic growth
- $b, \sigma$  Nemytskii operators, Lipschitz
- $(W_t)_{t \geq 0}$  cylindrical Wiener process
- $\Lambda \subset \mathbb{R}^d$  bounded domain
- control domain  $U$  non-convex.

# The Maximum Principle

# Pontryagin 1956: Controlled ODEs

Minimize

$$J(u) := \int_0^T l(x_t^u, u_t) dt + h(x_T^u)$$

subject to

$$\begin{cases} \partial_t x_t^u = b(x_t^u, u_t), & t \in [0, T] \\ x_0^u = x_0 \in \mathbb{R}^n. \end{cases}$$

Let  $\bar{u}$  be optimal. Then

$$\inf_{u \in U} H(t, \bar{x}_t, u) = H(t, \bar{x}_t, \bar{u}_t),$$

for almost every  $t \in [0, T]$ , where Hamiltonian  $H : [0, T] \times \mathbb{R}^n \times U \rightarrow \mathbb{R}$

$$H(t, x, u) := l(x, u) + \langle p_t, b(x, u) \rangle$$

where  $p_t$  is the adjoint state.

Minimize

$$J(u) := \mathbb{E} \left[ \int_0^T l(x_t^u, u_t) dt + h(x_T^u) \right]$$

subject to

$$\begin{cases} dx_t^u = b(x_t^u, u_t) dt + \sigma(x_t^u, u_t) dW_t, & t \in [0, T] \\ x_0^u = x_0 \in \mathbb{R}^n. \end{cases}$$

Let  $\bar{u}$  be optimal. Then

$$\inf_{u \in U} \mathcal{G}(t, \bar{x}_t, u) = \mathcal{G}(t, \bar{x}_t, \bar{u}_t),$$

for almost all  $(t, \omega) \in [0, T] \times \Omega$ , where generalized Hamiltonian

$$\mathcal{G} : [0, T] \times \mathbb{R}^n \times U \rightarrow \mathbb{R}$$

$$\mathcal{G}(t, x, u) := H(t, x, u) + \frac{1}{2} \text{tr}(\sigma(x, u)^* P_t \sigma(x, u)) + \text{tr}(\sigma(x, u)^* [q_t - P_t \sigma(\bar{x}_t, \bar{u}_t)])$$

where  $(p_t, q_t)$  first order adjoint state and  $P_t \in \mathbb{R}^{n \times n}$  second order adjoint state.

Minimize

$$J(u) := \mathbb{E} \left[ \int_0^T \int_{\Lambda} l(x_t^u(\lambda), u_t) d\lambda dt + \int_{\Lambda} h(x_T^u(\lambda)) d\lambda \right]$$

subject to

$$\begin{cases} dx_t^u = [\Delta x_t^u + b(x_t^u, u_t)] dt + \sigma(x_t^u, u_t) dW_t, & t \in [0, T] \\ x_0^u = x_0 \in L^2(\Lambda). \end{cases}$$

- Since Peng's result: Many generalizations to infinite-dimensional stochastic case by Du, Fuhrman, Frankowska, Guatteri, Hu, Li, Lü, Meng, Tang, Tessitore, Zhang, ...
- Major drawback in previous works: Restrictive assumptions on coefficients  $l$  and  $h$ , in particular excluding quadratic costs.
- Main obstacle: Characterization of second order adjoint state  $P_t \in L(H)$ .



# Spike Variation

Let  $\bar{u}$  be optimal. Fix  $\tau \in [0, T]$ ,  $\varepsilon > 0$ ,  $v \in U$ , and set

$$u_t^\varepsilon := \begin{cases} v, & \tau \leq t \leq \tau + \varepsilon, \\ \bar{u}_t & \text{otherwise.} \end{cases}$$

Then

$$0 \leq J(u^\varepsilon) - J(\bar{u}) = \mathbb{E} \left[ \int_0^T \int_\Lambda l(x_t^\varepsilon, u_t^\varepsilon) - l(\bar{x}_t, \bar{u}_t) d\lambda dt + \int_\Lambda h(x_T^\varepsilon) - h(\bar{x}_T) d\lambda \right].$$

Taylor expansions: For terminal costs we have

$$\mathbb{E} \left[ \int_\Lambda h(x_T^\varepsilon) - h(\bar{x}_T) d\lambda \right] \approx \mathbb{E} \left[ \int_\Lambda h_x(\bar{x}_T) y_T^\varepsilon d\lambda \right],$$

where  $y^\varepsilon$  satisfies linearized state equation

$$\begin{cases} dy_t^\varepsilon = [\Delta y_t^\varepsilon + b_x(\bar{x}_t, \bar{u}_t) y_t^\varepsilon + b(\bar{x}_t, u_t^\varepsilon) - b(\bar{x}_t, \bar{u}_t)] dt \\ \quad + [\sigma_x(\bar{x}_t, \bar{u}_t) y_t^\varepsilon + \sigma(\bar{x}_t, u_t^\varepsilon) - \sigma(\bar{x}_t, \bar{u}_t)] dW_t \\ y_0^\varepsilon = 0. \end{cases}$$

# Asymptotic of First Order Taylor Expansions

Applying Itô's formula to  $\|y_t^\varepsilon\|_{L^2(\Lambda)}^2$  yields

$$\begin{aligned}\|y_t^\varepsilon\|_{L^2(\Lambda)}^2 &= 2 \int_0^t \langle \Delta y_s^\varepsilon + b_x(\bar{x}_s, \bar{u}_s)y_s^\varepsilon + b(\bar{x}_s, u_s^\varepsilon) - b(\bar{x}_s, \bar{u}_s), y_s^\varepsilon \rangle ds \\ &\quad + 2 \int_0^t \langle y_s^\varepsilon, (\sigma_x(\bar{x}_s, \bar{u}_s)y_s^\varepsilon + \sigma(\bar{x}_s, u_s^\varepsilon) - \sigma(\bar{x}_s, \bar{u}_s)) dW_s \rangle \\ &\quad + \int_0^t \|\sigma_x(\bar{x}_s, \bar{u}_s)y_s^\varepsilon + \sigma(\bar{x}_s, u_s^\varepsilon) - \sigma(\bar{x}_s, \bar{u}_s)\|_{L_2(\Xi, L^2(\Lambda))}^2 ds.\end{aligned}$$

Itô correction term

$$\begin{aligned}&\int_0^t \|\sigma_x(\bar{x}_s, \bar{u}_s)y_s^\varepsilon + \sigma(\bar{x}_s, u_s^\varepsilon) - \sigma(\bar{x}_s, \bar{u}_s)\|_{L_2(\Xi, L^2(\Lambda))}^2 ds \\ &\leq 2 \int_0^t \|\sigma_x(\bar{x}_s, \bar{u}_s)y_s^\varepsilon\|_{L_2(\Xi, L^2(\Lambda))}^2 ds + 2 \int_\tau^{\tau+\varepsilon} \|\sigma(\bar{x}_s, v) - \sigma(\bar{x}_s, \bar{u}_s)\|_{L_2(\Xi, L^2(\Lambda))}^2 ds.\end{aligned}$$

The last term is of order  $\mathcal{O}(\varepsilon)$ ; we need  $o(\varepsilon)$ .  $\rightsquigarrow$  need second order Taylor expansions!

# Variational Equations

First order:

$$\begin{cases} dy_t^\varepsilon = [\Delta y_t^\varepsilon + b_x(\bar{x}_t, \bar{u}_t)y_t^\varepsilon + b(\bar{x}_t, u_t^\varepsilon) - b(\bar{x}_t, \bar{u}_t)] dt \\ \quad + [\sigma_x(\bar{x}_t, \bar{u}_t)y_t^\varepsilon + \sigma(\bar{x}_t, u_t^\varepsilon) - \sigma(\bar{x}_t, \bar{u}_t)] dW_t \\ y_0^\varepsilon = 0. \end{cases}$$

Second order:

$$\begin{cases} dz_t^\varepsilon = [\Delta z_t^\varepsilon + b_x(\bar{x}_t, \bar{u}_t)z_t^\varepsilon + \frac{1}{2}b_{xx}(\bar{x}_t, \bar{u}_t)y_t^\varepsilon y_t^\varepsilon + (b_x(\bar{x}_t, u_t^\varepsilon) - b_x(\bar{x}_t, \bar{u}_t))y_t^\varepsilon] dt \\ \quad + [\sigma_x(\bar{x}_t, \bar{u}_t)z_t^\varepsilon + \frac{1}{2}\sigma_{xx}(\bar{x}_t, \bar{u}_t)y_t^\varepsilon y_t^\varepsilon + (\sigma_x(\bar{x}_t, u_t^\varepsilon) - \sigma_x(\bar{x}_t, \bar{u}_t))y_t^\varepsilon] dW_t \\ z_0^\varepsilon = 0. \end{cases}$$

## Lemma

*It holds that*

$$\sup_{t \in [0, T]} \mathbb{E} \left[ \|y_t^\varepsilon\|_{L^2(\Lambda)}^2 \right] \leq C\varepsilon \quad \text{and} \quad \sup_{t \in [0, T]} \mathbb{E} \left[ \|z_t^\varepsilon\|_{L^2(\Lambda)}^2 \right] \leq C\varepsilon^2.$$

# Second Order Expansion of Cost Functional

From

$$0 \leq J(u^\varepsilon) - J(\bar{u}) = \mathbb{E} \left[ \int_0^T \int_\Lambda l(x_t^\varepsilon, u_t^\varepsilon) - l(\bar{x}_t, \bar{u}_t) d\lambda dt + \int_\Lambda h(x_T^\varepsilon) - h(\bar{x}_T) d\lambda \right]$$

we derive:

## Lemma

*It holds*

$$\begin{aligned} & \mathbb{E} \left[ \int_0^T \int_\Lambda l_x(\bar{x}_t(\lambda), \bar{u}_t)(y_t^\varepsilon(\lambda) + z_t^\varepsilon(\lambda)) + \frac{1}{2} l_{xx}(\bar{x}_t(\lambda), \bar{u}_t) y_t^\varepsilon(\lambda) y_t^\varepsilon(\lambda) d\lambda dt \right] \\ & + \mathbb{E} \left[ \int_\Lambda h_x(\bar{x}_T(\lambda))(y_T^\varepsilon(\lambda) + z_T^\varepsilon(\lambda)) + \frac{1}{2} h_{xx}(\bar{x}_T(\lambda)) y_T^\varepsilon(\lambda) y_T^\varepsilon(\lambda) d\lambda \right] \\ & + \mathbb{E} \left[ \int_0^T \int_\Lambda l(\bar{x}_t(\lambda), u_t^\varepsilon) - l(\bar{x}_t(\lambda), \bar{u}_t) d\lambda dt \right] \geq o(\varepsilon). \end{aligned}$$

Next step: Separate dependence on  $\varepsilon$ .

# First Order Adjoint State

Consider SPDE

$$\begin{cases} dy_t = [\Delta y_t + b_x(\bar{x}_t, \bar{u}_t)y_t + \varphi_t] dt + [\sigma_x(\bar{x}_t, \bar{u}_t)y_t + \psi_t] dW_t \\ y_0 = 0, \end{cases}$$

where  $(\varphi, \psi) \in L^2([0, T] \times \Omega; L^2(\Lambda)) \times L^2([0, T] \times \Omega; L_2(\Xi, L^2(\Lambda)))$ .

Construct linear functional

$$\mathcal{T}(\varphi, \psi) := \mathbb{E} \left[ \int_0^T \int_{\Lambda} l_x(\bar{x}_t(\lambda), \bar{u}_t)y_t(\lambda) d\lambda dt + \int_{\Lambda} h_x(\bar{x}_T(\lambda))y_T(\lambda) d\lambda \right].$$

By Riesz's representation theorem, there is a unique pair

$$(p, q) \in L^2([0, T] \times \Omega; L^2(\Lambda)) \times L^2([0, T] \times \Omega; L_2(\Xi, L^2(\Lambda)))$$

such that

$$\mathcal{T}(\varphi, \psi) = \mathbb{E} \left[ \int_0^T \langle \varphi_t, p_t \rangle_{L^2(\Lambda)} + \langle \psi_t, q_t \rangle_{L_2(\Xi, L^2(\Lambda))} dt \right]$$

for all  $(\varphi, \psi) \in L^2([0, T] \times \Omega; L^2(\Lambda)) \times L^2([0, T] \times \Omega; L_2(\Xi, L^2(\Lambda)))$ .

# First Order Adjoint Equation

Adjoint state property:

$$\mathbb{E} \left[ \int_0^T \langle y_t, l_x(\bar{x}_t, \bar{u}_t) \rangle dt + \langle h_x(\bar{x}_T), y_T \rangle \right] = \mathbb{E} \left[ \int_0^T \langle p_t, \varphi_t \rangle + \langle q_t, \psi_t \rangle dt \right].$$

Applying Itô's product rule yields

$$\begin{aligned} d\langle p_t, y_t \rangle_{L^2(\Lambda)} &= \langle p_t, dy_t \rangle_{L^2(\Lambda)} + \langle y_t, dp_t \rangle_{L^2(\Lambda)} + d\langle p, y \rangle_t \\ &\stackrel{!}{=} [\langle p_t, \varphi_t \rangle_{L^2(\Lambda)} + \langle q_t, \psi_t \rangle_{L_2(\Xi, L^2(\Lambda))} - \langle y_t, l_x(\bar{x}_t, \bar{u}_t) \rangle_{L^2(\Lambda)}] dt + dM_t. \end{aligned}$$

for some martingale  $(M_t)_{t \geq 0}$ . Thus,

$$\begin{cases} dp_t = - [\Delta p_t + b_x(\bar{x}_t, \bar{u}_t)p_t + l_x(\bar{x}_t, \bar{u}_t) + \langle \sigma_x(\bar{x}_t, \bar{u}_t), q_t \rangle_{L_2(\Xi, \mathbb{R})}] dt + q_t dW_t \\ p_T = h_x(\bar{x}_T). \end{cases}$$

Unique variational solution  $(p, q)$ , where

$$p \in L^2([0, T] \times \Omega; H_0^1(\Lambda)) \cap L^2(\Omega; C([0, T]; L^2(\Lambda)))$$

and

$$q \in L^2([0, T] \times \Omega; L_2(\Xi, L^2(\Lambda))).$$

# Backward Stochastic Differential Equations

Let  $(W_t)_{t \geq 0}$  be a Brownian motion and  $(\mathcal{F}_t)_{t \geq 0}$  its natural filtration. Consider the BSDE

$$\begin{cases} dp_t = 0 \\ p_T = \xi, \end{cases}$$

where  $\xi$  is  $\mathcal{F}_T$ -measurable.  $\triangleleft$  Adapted solution doesn't exist!

Natural candidate:  $p_t = \mathbb{E}[\xi | \mathcal{F}_t]$ . However, does not solve BSDE. Martingale representation theorem:

$$\mathbb{E}[\xi | \mathcal{F}_t] = \mathbb{E}[\xi] + \int_0^t q_s dW_s.$$

Restate original problem:

$$\begin{cases} dp_t = q_t dW_t \\ p_T = \xi. \end{cases}$$

For  $\xi = f(W_T)$ :

$$\mathbb{E}[\xi | \mathcal{F}_t] = \mathbb{E}[f(W_T) | \mathcal{F}_t] = S_{T-t} f(W_t) = S_T f(W_0) + \int_0^t \nabla S_{T-s} f(W_s) dW_s$$

Follow same route for quadratic terms

$$\mathbb{E} \left[ \int_0^T \int_{\Lambda} l_{xx}(\bar{x}_t(\lambda), \bar{u}_t) y_t^\varepsilon(\lambda) y_t^\varepsilon(\lambda) d\lambda dt + \int_{\Lambda} h_{xx}(\bar{x}_T(\lambda)) y_T^\varepsilon(\lambda) y_T^\varepsilon(\lambda) d\lambda \right].$$

Peng's idea in finite dimensions: Linearize using tensor product  $y_t^\varepsilon \otimes y_t^\varepsilon$  and derive equation on

$$\mathbb{R}^n \otimes \mathbb{R}^n \cong \mathbb{R}^{n \times n}.$$

Infinite dimensional analogue:

$$H \otimes H \cong L_2(H).$$

Problem: Quadratic costs require duality analysis between  $L_1(H)$  and  $L(H)$ .



Use explicit representation

$$\begin{aligned} L^2(\Lambda) \otimes L^2(\Lambda) &\cong L^2(\Lambda^2) \\ y \otimes z &\leftrightarrow ((\lambda, \mu) \mapsto y(\lambda)z(\mu)). \end{aligned}$$

Rewrite quadratic terms as

$$\begin{aligned} &\mathbb{E} \left[ \int_0^T \int_{\Lambda} l_{xx}(\bar{x}_t(\lambda), \bar{u}_t) y_t^\varepsilon(\lambda) y_t^\varepsilon(\lambda) d\lambda dt \right] \\ &= \mathbb{E} \left[ \int_0^T \int_{\Lambda} l_{xx}(\bar{x}_t(\lambda), \bar{u}_t) \delta(y_t^\varepsilon \otimes y_t^\varepsilon)(\lambda) d\lambda dt \right], \end{aligned}$$

where  $\delta : H_0^1(\Lambda^2) \rightarrow L^2(\Lambda)$  is defined by  $\delta(w)(\lambda) := w(\lambda, \lambda)$ . This expression is linear in  $y_t^\varepsilon \otimes y_t^\varepsilon$ .

# Second Order Adjoint Equation

## Theorem (Stannat, W., SICON 2021)

The equation

$$\left\{ \begin{array}{l} dP_t(\lambda, \mu) = -[\Delta P_t(\lambda, \mu) + (b_x(\bar{x}_t(\lambda), \bar{u}_t) + b_x(\bar{x}_t(\mu), \bar{u}_t))P_t(\lambda, \mu) \\ \quad + \langle \sigma_x(\bar{x}_t(\lambda), \bar{u}_t), \sigma_x(\bar{x}_t(\mu), \bar{u}_t) \rangle_{L_2(\Xi, \mathbb{R})} P_t(\lambda, \mu) \\ \quad + \langle \sigma_x(\bar{x}_t(\lambda), \bar{u}_t) + \sigma_x(\bar{x}_t(\mu), \bar{u}_t), Q_t(\lambda, \mu) \rangle_{L_2(\Xi, \mathbb{R})} \\ \quad + \delta^*(l_{xx}(\bar{x}_t(\lambda), \bar{u}_t)) + \delta^*(b_{xx}(\bar{x}_t(\lambda), \bar{u}_t)p_t(\lambda)) \\ \quad + \delta^*(\langle \sigma_{xx}(\bar{x}_t(\lambda), \bar{u}_t), q_t \rangle_{L_2(\Xi, \mathbb{R})})] dt + Q_t(\lambda, \mu) dW_t \\ P_T(\lambda, \mu) = \delta^*(h_{xx}(\bar{x}_T(\lambda))) \end{array} \right.$$

has a unique adapted solution  $(P, Q)$ , where

$$P \in L^2([0, T] \times \Omega; L^2(\Lambda^2)) \cap L^2(\Omega; C([0, T]; H^{-1}(\Lambda^2))),$$

and

$$Q \in L^2([0, T] \times \Omega; L_2(\Xi; H^{-1}(\Lambda^2))).$$

## Theorem (Stannat, W., SICON 2021)

Let  $(\bar{x}, \bar{u})$  be an optimal pair. Then there exist adapted processes

$$(p, q) \in L^2([0, T] \times \Omega; H_0^1(\Lambda)) \times L^2([0, T] \times \Omega; L_2(\Xi, L^2(\Lambda)))$$

satisfying the first order adjoint equation and adapted processes

$$(P, Q) \in L^2([0, T] \times \Omega; L^2(\Lambda^2)) \times L^2([0, T] \times \Omega; L_2(\Xi, H^{-1}(\Lambda^2)))$$

satisfying the second order adjoint equation such that

$$\inf_{u \in U} \mathcal{G}(t, \bar{x}_t, u) = \mathcal{G}(t, \bar{x}_t, \bar{u}_t),$$

for almost all  $(t, \omega) \in [0, T] \times \Omega$ , where  $\mathcal{G} : [0, T] \times L^2(\Lambda) \times U \rightarrow \mathbb{R}$

$$\begin{aligned} \mathcal{G}(t, x, u) := & \int_{\Lambda} l(x(\lambda), u) d\lambda + \langle p_t, b(x, u) \rangle_{L^2(\Lambda)} + \text{tr}(\sigma(x, u)^* q_t) \\ & + \frac{1}{2} \text{tr}(\sigma(x, u)^* P_t \sigma(x, u)) - \text{tr}(\sigma(x, u)^* P_t \sigma(\bar{x}_t, \bar{u}_t)). \end{aligned}$$

# The Dynamic Programming Approach

# Dynamic Programming Approach

Minimize

$$J(s, x; u) := \mathbb{E} \left[ \int_s^T \int_{\Lambda} l(x_t^u(\lambda), u_t) d\lambda dt + \int_{\Lambda} h(x_T^u(\lambda)) d\lambda \right]$$

over  $u \in \mathcal{U}_s$  subject to

$$\begin{cases} dx_t^u = [\Delta x_t^u + b(x_t^u, u_t)] dt + \sigma(x_t^u, u_t) dW_t, & t \in [s, T] \\ x_s^u = x \in L^2(\Lambda). \end{cases}$$

Introduce value function  $V : [0, T] \times L^2(\Lambda) \rightarrow \mathbb{R}$ ,

$$V(s, x) := \inf_{u \in \mathcal{U}_s} J(s, x; u).$$

Satisfies dynamic programming principle

$$V(s, x) = \inf_{u \in \mathcal{U}_s} \mathbb{E} \left[ \int_s^{\tau} l(x_t^u, u_t) dt + V(\tau, x_{\tau}^u) \right], \quad \forall \tau \in [s, T].$$

Can be used to derive optimality conditions.

Under smoothness assumptions, it holds

$$\begin{cases} V_s(t, \bar{x}_t) = -\langle \Delta \bar{x}_t, DV(t, \bar{x}_t) \rangle_{L^2(\Lambda)} - \mathcal{H}(\bar{x}_t, \bar{u}_t, DV(t, \bar{x}_t), D^2V(t, \bar{x}_t)) \\ DV(t, \bar{x}_t) = p_t \\ D^2V(t, \bar{x}_t)\sigma(\bar{x}_t, \bar{u}_t) = q_t, \end{cases}$$

where

$$\mathcal{H}(x, u, p, P) := \int_{\Lambda} l(x(\lambda), u) d\lambda + \langle p, b(x, u) \rangle_{L^2(\Lambda)} + \frac{1}{2} \text{tr}(\sigma(x, u)^* P \sigma(x, u)).$$

Generalizations dropping smoothness assumptions and using viscosity differentials up to first order in infinite dimensions by Cannarsa, Frankowska, Zhou, ...

# Viscosity Differentials

If  $V \in C^{1,2}([0, T] \times L^2(\Lambda))$ , it holds

$$\lim_{\tau \downarrow t, z \rightarrow x} \frac{1}{|\tau - t| + \|z - x\|^2} \left[ V(\tau, z) - V(t, x) - \partial_t V(t, x)(\tau - t) - \langle DV(t, x), z - x \rangle_{L^2(\Lambda)} - \frac{1}{2} \langle z - x, D^2 V(t, x)(z - x) \rangle_{L^2(\Lambda)} \right] = 0.$$

Weaker notion of differentiability:

## Definition (Viscosity Superdifferential)

We say  $(G, p, P) \in D_{t+, x}^{1,2,+} V(t, x)$  if

$$\limsup_{\tau \downarrow t, z \rightarrow x} \frac{1}{|\tau - t| + \|z - x\|^2} \left[ V(\tau, z) - V(t, x) - G(\tau - t) - \langle p, z - x \rangle_{L^2(\Lambda)} - \frac{1}{2} \langle z - x, P(z - x) \rangle_{L^2(\Lambda)} \right] \leq 0.$$

## Theorem (Stannat, W. (2022+))

For almost every  $t \in [0, T]$ , it holds

$$[-\langle \Delta \bar{x}_t, p_t \rangle_{H^{-1}(\Lambda) \times H_0^1(\Lambda)} - \mathcal{G}(t, \bar{x}_t, \bar{u}_t), \infty) \times \{p_t\} \times [P_t, \infty) \subset D_{t+,x}^{1,2,+} V(t, \bar{x}_t)$$

$\mathbb{P}$ -almost surely.

## Corollary (Stannat, W. (2022+))

It holds for almost all  $t \in [s, T]$

$$\mathcal{G}(t, \bar{x}_t, \bar{u}_t) \leq \mathcal{H}(t, \bar{x}_t, \bar{u}_t, p_t, P_t),$$

$\mathbb{P}$ -almost surely, i.e.,

$$\text{tr}(\sigma(\bar{x}_t, \bar{u}_t)(q_t - P_t \sigma(\bar{x}_t, \bar{u}_t))) \leq 0.$$



Value function formally satisfies Hamilton-Jacobi-Bellman equation

$$\begin{cases} V_s + \langle \Delta x, DV \rangle_{L^2(\Lambda)} + \inf_{u \in U} \mathcal{H}(x, u, DV, D^2V) = 0, & (s, x) \in [0, T] \times L^2(\Lambda) \\ V(T, x) = \int_{\Lambda} h(x(\lambda)) d\lambda, & x \in L^2(\Lambda) \end{cases}$$

where

$$\mathcal{H}(x, u, p, P) := \int_{\Lambda} l(x(\lambda), u) d\lambda + \langle p, b(x, u) \rangle_{L^2(\Lambda)} + \frac{1}{2} \text{tr}(\sigma(x, u)^* P \sigma(x, u)).$$

## Definition (Viscosity Solution, Bounded Case)

$V$  is viscosity subsolution, if

- $V(T, x) \leq \int_{\Lambda} h(x(\lambda)) d\lambda, \quad x \in L^2(\Lambda);$
- for every  $(G, p, P) \in D_{t,x}^{1,2,+} V(t, x)$

$$G + \langle Ax, p \rangle + \inf_{u \in U} \mathcal{H}(x, u, p, P) \geq 0.$$

# Viscosity Solutions II

It holds:

$$(G, p, P) \in D_{t,x}^{1,2,+} v(t, x)$$



$\exists \phi \in C^{1,2}((s, T) \times L^2(\Lambda))$  such that:

- 1  $v - \phi$  attains maximum at  $(t, x)$ ,
- 2  $(\phi(t, x), \partial_t \phi(t, x), D\phi(t, x), D^2\phi(t, x)) = (v(t, x), G, p, P)$ .

Equivalent definition of viscosity solution in the bounded case (!):

## Definition (Viscosity Solution, Bounded Case)

$V$  is a viscosity subsolution, if

- $V(T, x) \leq \int_{\Lambda} h(x(\lambda)) d\lambda, \quad x \in L^2(\Lambda);$
- $\forall \phi \in C^{1,2}((s, T) \times L^2(\Lambda))$  such that  $V - \phi$  attains maximum at  $(t, x)$ , it holds

$$\phi_s(t, x) + \langle Ax, D\phi(t, x) \rangle + \inf_{u \in U} \mathcal{H}(x, u, D\phi(t, x), D^2\phi(t, x)) \geq 0.$$

## Corollary (Stannat, W. (2022+))

It holds for almost all  $t \in [s, T]$

$$\mathcal{G}(t, \bar{x}_t, \bar{u}_t) \leq \mathcal{H}(t, \bar{x}_t, \bar{u}_t, p_t, P_t).$$

Proof (bounded case!): Take test function  $\phi$  satisfying

$$(\phi(t, \bar{x}_t), \phi_t(t, \bar{x}_t), D\phi(t, \bar{x}_t), D^2\phi(t, \bar{x}_t)) = (V(t, \bar{x}_t), -\langle A\bar{x}_t, p_t \rangle - \mathcal{G}(t, \bar{x}_t, \bar{u}_t), p_t, P_t).$$

Since  $V$  is a viscosity solution of the HJB equation, we have

$$\begin{aligned} 0 &\leq \phi_t(t, \bar{x}_t) + \langle A\bar{x}_t, D\phi(t, \bar{x}_t) \rangle + \inf_{u \in U} \mathcal{H}(t, \bar{x}_t, u, D\phi(t, \bar{x}_t), D^2\phi(t, \bar{x}_t)) \\ &= -\mathcal{G}(t, \bar{x}_t, \bar{u}_t) - \langle A\bar{x}_t, p_t \rangle + \langle A\bar{x}_t, p_t \rangle + \inf_{u \in U} \mathcal{H}(t, \bar{x}_t, u, p_t, P_t) \\ &\leq -\mathcal{G}(t, \bar{x}_t, \bar{u}_t) + \mathcal{H}(t, \bar{x}_t, \bar{u}_t, p_t, P_t) \end{aligned}$$



# Generalized Hamiltonian vs. Hamiltonian

HJB equation:

$$\begin{cases} V_s + \langle \Delta x, DV \rangle_{L^2(\Lambda)} + \inf_{u \in U} \mathcal{H}(x, u, DV, D^2V) = 0, & (s, x) \in [0, T] \times L^2(\Lambda) \\ V(T, x) = \int_{\Lambda} h(x(\lambda)) d\lambda, & x \in L^2(\Lambda) \end{cases}$$

Problematic term:

$$\langle \Delta x, D\phi(t, x) \rangle_{L^2(\Lambda)}, \quad x \in L^2(\Lambda).$$

$\rightsquigarrow$  Need to restrict class of test functions.

We only need to make sense of

$$\langle \Delta \bar{x}_t, D\phi(t, \bar{x}_t) \rangle_{L^2(\Lambda)},$$

where  $\bar{x}_t \in H_0^1(\Lambda)$ ,  $dt \otimes \mathbb{P}$ -a.s.

# Verification Theorem

Minimize

$$J(s, x; u) := \mathbb{E} \left[ \int_s^T L(x_t^u, u_t) dt + H(x_T^u) \right]$$

over  $u \in \mathcal{U}_s$  subject to

$$\begin{cases} dx_t^u = [\Delta x_t^u + B(x_t^u, u_t)] dt + \Sigma(x_t^u, u_t) dW_t, & t \in [s, T] \\ x_s^u = x \in L^2(\Lambda). \end{cases}$$

Smooth verification theorem: Let  $(x^*, u^*)$  be admissible such that

$$u_t^* \in \arg \min_{u \in U} \mathcal{H}(x_t^*, u, DV(t, x_t^*), D^2V(t, x_t^*)),$$

for almost every  $(t, \omega)$ . Then  $(x^*, u^*)$  is optimal.

Verification theorems in the stochastic case in the framework of viscosity solutions by Fabbri, Gozzi, Li, Świąch, Yong, Zhou, ...

# Verification Theorem

Minimize

$$J(s, x; u) := \mathbb{E} \left[ \int_s^T L(x_t^u, u_t) dt + H(x_T^u) \right]$$

over  $u \in \mathcal{U}_s$  subject to

$$\begin{cases} dx_t^u = [\Delta x_t^u + B(x_t^u, u_t)] dt + \Sigma(x_t^u, u_t) dW_t, & t \in [s, T] \\ x_s^u = x \in L^2(\Lambda). \end{cases}$$

Smooth verification theorem: Let  $(x^*, u^*)$  be admissible such that

$$V_s(t, x_t^*) + \langle \Delta x_t^*, DV(t, x_t^*) \rangle_{L^2(\Lambda)} + \mathcal{H}(x_t^*, u_t^*, DV(t, x_t^*), D^2V(t, x_t^*)) = 0$$

for almost every  $(t, \omega)$ . Then  $(x^*, u^*)$  is optimal.

Verification theorems in the stochastic case in the framework of viscosity solutions by Fabbri, Gozzi, Li, Świąch, Yong, Zhou, ...

## Theorem (Stannat, W. (2022+))

Assume

- $\|\Sigma(x, u)\|_{L_2(\Xi, H_0^1(\Lambda))} \leq C(1 + \|x\|_{H_0^1(\Lambda)})$
- $V(t + \tau, x) - V(t, x) \leq C(1 + \|x\|_{H_0^1(\Lambda)}^2)\tau$
- $V(t, \cdot) - C\|\cdot\|_{L^2(\Lambda)}^2$  is concave.

Let  $(x^*, u^*)$  be an admissible pair. Suppose there are adapted processes  $(G, p, P)$  taking values in  $\mathbb{R}$ ,  $H_0^1(\Lambda)$  and  $L_2(L^2(\Lambda))$ , such that for almost all  $t \in [s, T]$ :

$$(G_t, p_t, P_t) \in D_{t+, x}^{1,2,+} V(t, x_t^*)$$

$\mathbb{P}$ -almost surely, and

$$\mathbb{E} \left[ \int_s^T G_t + \langle \Delta x_t^*, p_t \rangle_{H^{-1}(\Lambda) \times H_0^1(\Lambda)} + \mathcal{H}(x_t^*, u_t^*, p_t, P_t) dt \right] \geq 0.$$

Then  $(x^*, u^*)$  is an optimal pair.



W. Stannat, L. Wessels

Peng's maximum principle for stochastic partial differential equations  
SIAM J. Control Optim. 59 (2021), pp. 3552–3573.



W. Stannat, L. Wessels

Necessary and sufficient conditions for optimal control of semilinear SPDEs  
Submitted, arXiv:2112.09639.



W. Stannat, L. Wessels

Deterministic control of stochastic reaction-diffusion equations  
Evol. Equ. Control Theory 10 (2021), pp. 701–722.



S. Peng

A general stochastic maximum principle for optimal control problems  
SIAM J. Control Optim. 28 (1990), pp. 966–979.



L. S. Pontryagin, V. G. Boltyanskii, R. V. Gamkrelidze, and E. F. Mishchenko

The mathematical theory of optimal processes  
Publishers John Wiley & Sons, Inc., New York, 1962.